

# *Herniaria ciliolata* ssp *ciliolata*

Key:

**Dots** refer to the native sites of the species

**Numbers** refer to the nearby Botanical Collections



## *Starting references*

### **Family**

Caryophyllaceae

### **IUCN category (2001)**

Vulnerable. Endemic.

### **Habit**

Mat-forming perennial herb.

### **Habitat**

Short, open vegetation, usually on dry, base-rich soils or rocks on S-facing slopes; on coastal cliff-slopes, valley sides, dune grassland, rock outcrops and open patches in heathlands. Lowland.

### **Reasons for decline**

### **Distribution in wild**

| Country | Locality & Vice County               | Sites<br>(10km <sup>2</sup> occurrences) | Population<br>(plants) |
|---------|--------------------------------------|--|------------------------|
| England | W Cornwall                           | 2  |                        |
|         | Guernsey & Alderney, Channel Islands | 2  |                        |

### *Ex situ Collections*

#### **Gardens close to the region of distribution of the species**

- 1 Duchy College
- 2 St Michael's Mount (NT)
- 3 Trebah Garden
- 4 Glendurgan Gardens (NT)
- 5 Trellisick (NT)
- 6 Tregothnan Botanic Garden
- 7 Eden Project

#### **Gardens with specialisation on family Caryophyllaceae**

University of Oxford Botanic Garden

#### **Potential to grow the species in *ex situ* Collections**

### *Conservation information*

#### **Linkages to BAPs**

Unknown

#### **Habitat Management**

Unknown

#### **Conservation programmes**

Unknown

### **Web References**

- Images: [http://www.ukwildflowers.com/Web\\_pages/herniaria\\_ciliolata\\_fringed\\_rupturewort.htm](http://www.ukwildflowers.com/Web_pages/herniaria_ciliolata_fringed_rupturewort.htm)